

## Angiotensin Converting Enzyme 2 (ACE2) Polyclonal Antibody

### ORDERING INFORMATION

**Catalog No.:** 19504

**Size:** 100ul (1mg/ml)

**Format:** Antigen affinity-purified antibody in PBS, 0.02% sodium azide.

### BACKGROUND

Angiotensin-converting enzyme 2 (ACE2) is a zinc containing metalloenzyme located on the surface of endothelial and other cells in the lungs, arteries, heart, kidney, and intestines. The primary function of ACE2 is to offset activity of angiotensin-converting enzyme (ACE). ACE cleaves angiotensin I into the vasoconstrictor angiotensin II. ACE2 in turn cleaves angiotensin II into the vasodilator angiotensin 1–7. In addition, ACE2 is the main entry point into cells for some coronaviruses including HCoV-NL63, SARS-CoV (the coronavirus that causes SARS) and SARS-CoV-2 (the coronavirus that causes COVID-19). More precisely, the binding of the spike S1 protein of SARS-CoV and SARS-CoV2 to the enzymatic domain of ACE2 on the surface of cells results in endocytosis and translocation of both the virus and the enzyme into endosomes located within cells.

### SPECIFICATION SUMMARY

**Antigen:** Synthetic peptide corresponding to amino acids near the N-terminus of human ACE2 protein.

**Accession no.:** Q9BYF1


**Gene ID:** 59272

**Host Species:** Rabbit

**Specificity:** Human ACE2. Does not cross-react with ACE1.


### APPLICATION

**Immunoblotting:** use at 1ug/ml. A band of ~92kDa is detected.



Detection of ACE2 in human kidney lysate with #19504 at 1ug/ml.

**IHC:** use at 2ug/ml.



Detection of ACE2 in paraffin-embedded human kidney with #19504 at 2ug/ml.

These are recommended concentrations. Endusers should determine optimal concentrations for their applications

### STORAGE AND STABILITY

This antibody is stable for at least one (1) year at -20°C. Store product in appropriate aliquots to avoid multiple freeze-thaw cycles.

*For in vitro investigational use only. Not for use in therapeutic or diagnostic procedures.*